

Calibrated Fibrations on Complete Manifolds via Torus Action

Edward Goldstein

February 1, 2008

Abstract

In this paper we will investigate torus actions on complete manifolds with calibrations. For Calabi-Yau manifolds M^{2n} with a Hamiltonian structure-preserving k -torus action we show that any smooth symplectic reduction has a natural holomorphic volume form. Moreover Special Lagrangian (SLag) submanifolds of the reduction lift to SLag submanifolds of M , invariant under the torus action. If $k = n - 1$ and $H^1(M, \mathbb{R}) = 0$ then we prove that M is a fibration with generic fiber being a SLag submanifold. As an application we will see that crepant resolutions of singularities of a finite Abelian subgroup of $SU(n)$ acting on \mathbb{C}^n have SLag fibrations. We study SLag submanifolds on the total space $K(N)$ of the canonical bundle of a Kahler-Einstein manifold N with positive scalar curvature and give a conjecture about fibration of $K(N)$ by SLag sub-varieties, which we prove if N is toric. We will also get somewhat weaker results for coassociative submanifolds of a G_2 -manifold M^7 , which admits a 3-torus, a 2-torus or an $SO(3)$ action.

1 Introduction

In this paper we will use structure preserving torus actions on non-compact manifolds with calibrations to construct calibrated submanifolds (both for Calabi-Yau manifolds and for 7-manifolds with a G_2 structure). We will assume that no element of the torus acts trivially. Throughout the paper we will use the notion of a calibrated fibration:

Definition 1 *Let (M, φ) be a Riemannian manifold with a calibrating form φ . Then we say that M has a calibrated fibration on it if there is a surjective map $\alpha : M \mapsto V$ onto a topological space V and a subset $S \subset M$ s.t.*

- i) For any point $m \in M - S$ the level set L_m of α through m is a smooth submanifold, calibrated by φ .*
- ii) The set S is locally contained in a finite union of submanifolds of codimension ≥ 4 in M .*

In Section 2 we consider a Kahler manifold (M^{2n}, ω) with a non-vanishing holomorphic $(n, 0)$ form φ . We can define, as in [4], Special Lagrangian (SLag) submanifolds L by the conditions:

$$\omega|_L = 0, \quad \text{Im}\varphi|_L = 0.$$

If g is the Riemannian metric corresponding to ω , then we can conformally scale g to a metric g' on M so that the form φ will have length $\sqrt{2}^n$ with respect to g' . Then SLag submanifolds will be calibrated by $\text{Re}\varphi$ with respect to g' . In particular, they will be minimal submanifolds of (M, g') and Lagrangian submanifolds of ω .

If M is compact and simply connected, then for any Kahler form ω on M Yau's celebrated resolution of the Calabi conjecture gives a (unique) Ricci-flat Kahler form ω' in the same cohomology class as ω (see [15]). Also the SYZ conjecture (see [13]) states that $(M, \text{Re}\varphi)$ has a calibrated fibration with generic fiber being a SLag torus with respect to a Ricci-flat Kahler metric. We can ask an analogous question for any Kahler metric on M and we showed in [4] that this holds for a choice of Kahler metric on a Borcea-Voisin threefold. In this paper we will be interested in non-compact Calabi-Yau manifolds with a structure-preserving torus action. The main results of Section 2 are as follows:

Theorem 1 *Suppose we have a Hamiltonian structure-preserving T^k -action on M^{2n} . Then any smooth symplectic reduction M_{red} has a natural holomorphic volume form φ_{red} . Moreover SLag submanifolds of $M_{\text{red}}, \omega_{\text{red}}, \varphi_{\text{red}}$ lift to SLag submanifolds of M , invariant under T^k -action. Vice versa, let L be a connected, T -invariant SLag submanifold of M s.t. T acts freely on L . Then L lies on a level set a of the moment map μ and one can find a SLag submanifold L' in the smooth part of the symplectic reduction through level set a s.t. L' lifts to L .*

Theorem 2 *Suppose that $k = n - 1$ and $H^1(M, \mathbb{R}) = 0$. Then M has a calibrated fibration α over an open subset of \mathbb{R}^n with the set S of singular points being the non-regular points of the torus action (i.e points where the differential of the action is not injective). Moreover for a generic point p (outside of a countable union of $(n - 2)$ -dimensional planes in \mathbb{R}^n), the fiber $\alpha^{-1}(p)$ is a smooth SLag submanifold. Connected components of each smooth fiber are diffeomorphic to an $(n - 1)$ -torus times \mathbb{R} . Singular fibers have singularities of codimension at least 2, and near singular points they are diffeomorphic to a product of a cone with a Euclidean ball.*

If we make certain assumptions on the set of non-regular points of T -action, then we can replace the countable union by a finite union in Theorem 2 (see Theorem 3 in Section 2).

In Section 3 we consider a 7-manifold M with a G_2 -form φ (see [3]). Let $H^1(M, \mathbb{R}) = 0$. If M has a 3-torus action, then M is covered by a family of non-intersecting coassociative submanifolds. Suppose M admits a 2-torus action. Then we will define certain G_2 -reductions M_{red} , which are symplectic

4-manifolds with a compatible almost complex structure and trivial canonical bundle. We will see that 2-dimensional complex sub-varieties of M_{red} lift to T -invariant, coassociative submanifolds of M . Finally suppose M admits an $SO(3)$ -action, which is not regular in at least one point. If $SO(3)$ acts freely on the set M' of regular points of the $SO(3)$ -action, then M' is covered by a family of non-intersecting coassociative submanifolds, diffeomorphic to $SO(3) \times \mathbb{R}$.

In Section 4 we will consider some applications of the results of the two previous sections. For the Calabi-Yau case, we will show that crepant resolutions of singularities of a finite Abelian subgroup of $SU(n)$ acting on \mathbb{C}^n have SLag fibrations. Also for any Kahler-Einstein manifold N with positive scalar curvature, the total space $K(N)$ of its canonical bundle is a Calabi-Yau manifold (see [12]). We investigate SLag submanifolds on $K(N)$. For each orientable minimal Lagrangian submanifold of N we associate a 1-parameter family $(L_\lambda | \lambda \in \mathbb{R})$ of SLag submanifolds of $K(N)$. Also L_0 is invariant under scaling of $K(N)$ by a real number. For any compact Kahler manifold N^{2n} with an effective n -torus action we prove that one of the regular orbits of the action is a minimal Lagrangian submanifold of N . If N is Kahler-Einstein with nonzero scalar curvature t and toric, then we prove that precisely 1 such orbit L is a minimal Lagrangian submanifold of N . For $t > 0$ we use Theorem 2 to construct a SLag fibration on $K(N)$ and we prove that all fibers are asymptotic at infinity to the fiber L_0 . We conjecture that any K-E manifold N with positive scalar curvature has a minimal Lagrangian submanifold L . Moreover $K(N)$ fibers with generic fiber being a SLag submanifold of $K(N)$ and all fibers are asymptotic to L_0 at infinity.

In the G_2 case, Bryant and Salamon have constructed in [3] some examples of complete metrics with holonomy G_2 . Some metrics are on the total space of the spin bundle over a 3-dimensional space form. Others are on a total space of a bundle Λ^2_- of anti-self-dual 2-forms over a self-dual Einstein 4-manifold. Many examples admit T^2 and $SO(3)$ -actions, and we show that in one example the G_2 manifold M can be covered by non-intersecting coassociative $SO(3)$ -invariant submanifolds.

Acknowledgments : This paper is a part of author's work towards his Ph.D. at the Massachusetts Institute of Technology. The author wants to express his gratitude to his advisor, Tom Mrowka, for initiating him into this subject and for continuing support.

After writing this paper the author learned that Mark Gross has independently obtained results, which are similar to some results of Sections 2 and 4.3.

2 Torus action on Calabi-Yau manifolds

Let $(M^{2n}, \omega, \varphi)$ be a Calabi-Yau manifold with a structure-preserving Hamiltonian T^k -action. For any element v of the Lie algebra \mathcal{G} of T^k we associate the infinitesimal flow vector field X_v on M , induced by the differential of the action. Then the X_v commute and their flows preserve ω and φ . Let v_1, \dots, v_l

be elements of \mathcal{G} and X_1, \dots, X_l be the corresponding vector fields on M . We claim that the $(n-l, 0)$ -form $\varphi' = i_{X_1} \dots i_{X_l} \varphi$ (obtained by contraction of φ by the vector fields X_1, \dots, X_l) is a closed $(n-l, 0)$ -form. We will prove it by induction on l . If $l = 1$ then the X_1 -flow preserves φ and so

$$0 = \mathcal{L}_{X_1} \varphi = d(i_{X_1} \varphi) = d\varphi'$$

Now we use induction on l . The X_1 flow preserves X_2, \dots, X_l and it preserves φ , so it preserves $\varphi'' = i_{X_2} \dots i_{X_l} \varphi$. So

$$0 = \mathcal{L}_{X_1} \varphi'' = d(i_{X_1} \varphi'') + i_{X_1} (d\varphi'')$$

and we are done by induction.

A moment map μ for the T^k -action is a map $\mu : M \mapsto \mathcal{G}^*$ (the dual Lie Algebra of T^k), which satisfies

$$d(\mu(v)) = i_{X_v} \omega$$

for any $v \in \mathcal{G}$. The moment map is T^k -invariant.

Theorem 1 *Any smooth symplectic reduction M_{red} has a natural holomorphic volume form φ_{red} . Moreover SLag submanifolds of M_{red} lift to SLag submanifolds of M , invariant under T^k -action. Vice versa let L be a connected, T -invariant SLag submanifold of M . Then L lies on a level set a of the moment map μ and moreover there is a SLag submanifold L' on a smooth part of symplectic reduction through level set a s.t. L is the lift of L' .*

Proof : Consider a level set Σ_a of the moment map μ , on which T^k acts freely. Since μ is T^k -invariant, the vector fields X_v are tangent to Σ_a . Consider the bundle $V = \text{span}(X_v)$ over Σ_a . Since $d(\mu(v)) = i_{X_v} \omega$, the tangent bundle to Σ_a is the ω -orthogonal complement of V . Also $V \cap JV = 0$ (here J is the complex structure on M). Let $W = (V \oplus JV)^\perp$ (here \perp is with respect to the metric). Then W is a complex vector bundle over Σ_a , the tangent bundle to Σ_a is $W \oplus V$ and the quotient of W by the T -action can be viewed as a tangent bundle to the symplectic reduction $M_{red} = \Sigma_a / T^k$.

Let v_1, \dots, v_k be a basis for the Lie algebra of T^k and X_1, \dots, X_k corresponding vector fields on M . Let $\varphi' = i_{X_1} \dots i_{X_k} \varphi$. Then as we have seen, φ' is a holomorphic $(n-k, 0)$ form on M . Also $\varphi'|_W$ is a holomorphic volume form on W . Since φ' is T^k -invariant, it is clear that there is a unique $(n-k, 0)$ form φ_{red} on M_{red} s.t. $\pi^*(\varphi_{red}) = \varphi'$ on Σ_a (here $\pi : \Sigma_a \mapsto M_{red}$ is the quotient map). Now φ_{red} is closed (since φ' is), and hence it is a holomorphic volume form on M_{red} .

Let L' be a SLag submanifold of M_{red} and $L = \pi^{-1}(L')$. It is clear that L is a SLag submanifold of M , invariant under T^k -action. Vice versa, let L be a connected SLag submanifold of M , invariant under T^k -action and T acts freely on L . Since L is invariant under the torus action, the vector fields X_v are tangent to L . Let u be some vector, tangent to L . Since L is Lagrangian, we have

$$0 = \omega(X_v, u) = d(\mu(v))(u)$$

So the differential of the moment map is zero on L , hence L lies on some level set Σ_a of the moment map. Since T -action is free on L , it is freely on a neighbourhood U of L in Σ_a . Moreover U is a smooth submanifold of M . So $U_{red} = U/T$ will be an open set in the smooth part of the symplectic reduction through a . It is clear that the quotient $L' = L/T^k$ is a SLag submanifold on U_{red} . Q.E.D.

In case $k = n - 1$ M has a calibrated fibration by SLag submanifolds, invariant under T^k -action:

Theorem 2 *Let $k = n - 1$ and $H^1(M, \mathbb{R}) = 0$. Then*

i) M has a calibrated fibration α over an open subset of \mathbb{R}^n with the set S of singular points being the non-regular points of the T -action.

ii) For a generic point p (outside of a countable union of $(n - 2)$ -planes in \mathbb{R}^n), the fiber $\alpha^{-1}(p)$ is a smooth SLag submanifold of M .

iii) Connected components of each smooth fiber are diffeomorphic to a product of an $(n - 1)$ -torus with \mathbb{R} .

iv) Singular fibers have singularities of codimension at least 2, and near a singular point they are diffeomorphic to a product of a cone with a Euclidean ball.

Proof: Define the form φ' as in the proof of Theorem 1. Then φ' is a holomorphic $(1, 0)$ -form, invariant under the torus action. Since $H^1(M, \mathbb{R}) = 0$, there is a holomorphic function $f = \eta + i\xi$ s.t. $df = \varphi'$. It is clear that f is also invariant under T^k -action. Let L be a connected SLag submanifold, invariant under the torus action. As we have seen, L must lie on the level set of the moment map μ . Also, since L is Special, one easily deduces that $Im\varphi'|_L = 0$, so L must lie on a level set of $\xi = Imf$, i.e. L lies on a level set of n -functions

$$\mu = a, \quad \xi = c$$

The moment map μ goes to \mathcal{G}^* , which we identify with \mathbb{R}^{n-1} by choosing a basis of \mathcal{G} . We define $\alpha = (\mu, \xi) : M \mapsto \mathbb{R}^n$.

Let S be the set of non-regular points of the torus action. We claim that a level set L_m of (μ, ξ) , that passes through a regular point $m \in M - S$ is smooth n -dimensional SLag submanifold of M near m . Indeed let Σ_a be the level set of the moment map passing through m and V and W be vector bundles on Σ_a near m as in the proof of Theorem 1. Let v_1, \dots, v_k be a basis for the Lie algebra of T^k . Then $d\mu(v_1), \dots, d\mu(v_k)$ is basis of $(JV)^*$. Also those 1-forms vanish on W . Now $d\xi = Im\varphi'$ restricted to W is non-zero. So the forms $d\mu(v_1), \dots, d\mu(v_k), d\xi$ are linearly independent at m , and so the level set L_m is a smooth submanifold of M near m .

Next we prove that L_m is SLag near m . Since ξ and μ are T^k -invariant, then so is L_m . So the X_v are in the tangent space to L_m at m . Since L_m is on the level set of μ , the tangent space to L_m at m is ω -orthogonal to X_v 's, so it must be Lagrangian. Also $Im\varphi'|_{L_m} = 0$ implies that L_m is Special at m .

Now we prove **iii)**: Let L be a level set of (μ, ξ) s.t. all points on L are regular points for T -action on M . Let L' be a connected component of L . We

have $Re\varphi' = d\eta$. One easily sees that $Re\varphi'|_L \neq 0$ for all points of L . One also easily shows that $\nabla\eta$ along L is tangent to L , hence it coincides with $\nabla(\eta|_L)$.

Consider a T^{n-1} -orbit T on L' . T lives on some level set of η on L' , hence it must coincide with a connected component of this level set. The normalized gradient flow of T by $\frac{\nabla\eta}{|\nabla\eta|^2}$ on L' gives a diffeomorphism between L' and $T^{(n-1)}$ times \mathbb{R} . Indeed this flow is defined on an interval (a_-, a_+) , there a_-, a_+ are independent on the choice of a point in T . One easily deduces that the orbit of T under this flow is isolated in L , hence it is equal to L' .

Next we prove **ii)** and show that S is locally contained in a finite union of submanifolds of codimension 4 in M . To prove this we need to understand the picture near a point in S . Let $m \in S$. The differential of the action is not injective at m and m has a stabilizer T' of positive dimension l and an orbit O . To prove **ii)** we need to see what is the image of (μ, ξ) on S near m .

The symplectic form ω restricts trivially to O , hence we have $\omega = d\gamma$ for some 1-form γ in a neighbourhood of O . Since ω is T^{n-1} -invariant, we can make γ invariant as well (by integrating over T^{n-1}). For any $v \in \mathcal{G}$ we have $0 = \mathcal{L}_{X_v}\gamma = d(\gamma(X_v)) + i_{X_v}\omega$. So the map $\mu'(v) = -\gamma(X_v)$ is a moment map near O and $\mu - \mu'$ is a constant. Obviously $Im\varphi|_O = 0$, so we can write $Im\varphi = d\beta$ for a T -invariant $(n-1)$ -form β in a neighbourhood of O . Let $\xi' = \beta(X_1, \dots, X_{n-1})$. Arguing as in the proof of Theorem 1 we get that $\varphi' = d\xi'$, so $\xi - \xi'$ is a constant (here $\xi = Imf$ as before). We will prove that the image of S near O by (μ', ξ') is contained in a finite union of $(n-2)$ -planes in \mathbb{R}^n . This will prove that one can find a neighbourhood U of m and a finite union H of $(n-2)$ -planes in \mathbb{R}^n s.t. $(\mu, \xi)(S \cap U) \subset H$. Since M is paracompact, one can find a countable union of $(n-2)$ -planes H' in \mathbb{R}^n s.t. $(\mu, \xi)(S) \subset H'$ and **ii)** follows.

Obviously $\xi' = \beta(X_1, \dots, X_{n-1}) = 0$ on S . Next we prove that there is a collection v_1, \dots, v_l of linearly independent elements of \mathcal{G} s.t. at any point $p' \in S$ the flow field corresponding to $v_i - v_j$ vanishes for some i and j . This will imply that the image of μ' on S will lie on a finite collection of hyper-planes in the dual Lie Algebra \mathcal{G}^* . Hence the image of (μ', ξ') on S near O will be contained in a finite collection of $(n-2)$ -planes.

We have the tangent bundle TO and the normal bundle $N(O)$, which splits as a direct sum $N(O) = J(TO) \oplus W$, $W = (TO + J(TO))^\perp$. W is a complex vector bundle of dimension $l+1$. T' -action preserves O and it's differential preserves TO and $J(TO)$, hence T' acts faithfully on W (because no element of T' acts trivially on M). We get an injective homomorphism ρ from T' to $SU(W)$, whose image is in some maximal torus of $SU(W)$. By dimension count this image is the maximal torus of $SU(W)$. We identify a small neighbourhood V of O in M with a small ball in $N(O)$ by the exponential map. Then the action of T' under this identification is trivial on $J(TO)$ and equal to the action by representation ρ on W . Let $\pi : V \simeq N(O) \rightarrow O$ be the projection. Take any element v of the Lie Algebra of T . Then it is clear that $\pi_*(X_v) = 0$ iff v is in the Lie Algebra \mathcal{G}' of T' . So a point $p \in V$ is in S iff there is an element $0 \neq v \in \mathcal{G}'$ s.t. $X_v(p) = 0$.

We identify the fiber $W(m)$ at m with \mathbb{C}^{l+1} . We have a torus $T' \simeq T^l$ acting as a maximal torus of $SU(l+1)$ on \mathbb{C}^{l+1} by

$$(e^{i\theta_1}, \dots, e^{i\theta_l})(z_1, \dots, z_{l+1}) = (e^{i\theta_1} z_1, \dots, e^{i\theta_l} z_l, e^{-\sum \theta_i} z_{l+1}) \quad (1)$$

It is clear that the non-regular points in $\pi^{-1}(m)$ are subspaces $H_{i,j} = (z_i = z_j = 0) \oplus J(TO)$ and the vector field $\partial_{\theta_i} - \partial_{\theta_j}$ is in the kernel of the differential of the action at these points. Let U' be some submanifold of T , which passes through $0 \in T$ and is transversal to T' at 0. Then the image of m under U' -action is a neighbourhood of m in O . Let $U_{i,j}$ be the image of $H_{i,j}$ under U' -action. The flow vector field of $\partial_{\theta_i} - \partial_{\theta_j}$ vanishes along $U_{i,j}$. We claim that near m S is contained in the union of $U_{i,j}$. Indeed let $p' \in S$ and let $m' = \pi(p')$. There is an element of U' which sends m to m' and hence it sends p to p' for some $p \in \pi^{-1}(m)$. One easily deduces that $p \in \bigcup H_{i,j}$, hence $p' \in \bigcup U_{i,j}$. Thus we can take $v_i = \partial_{\theta_i}$ and we are done. Moreover $U_{i,j}$ is obviously a submanifold of M of codimension 4, thus we also proved that S is locally contained in a finite union of submanifolds of codimension 4 in M .

To complete the proof of **i) and iv)** we still need to investigate the structure of the singular fiber L_m through m and to prove that the image of (μ, ξ) is open. Let e_1, \dots, e_l be the basis of Lie algebra of T' . We extend it by e_{l+1}, \dots, e_{n-1} to be the basis of \mathcal{G} . Let $\mu'' = (\mu(e_{l+1}), \dots, \mu(e_{n-1}))$. Then the differential of μ'' is surjective along O and the level set Σ of μ'' through m is a smooth submanifold of M (containing O). Also obviously $L_m \subset \Sigma$. We can investigate L_m by means of a local symplectic reduction. Let $Q = \text{span}(e_{l+1}, \dots, e_{n-1})$. Take a small ball U containing the origin in Q . U can be identified with a submanifold (still called U) of T^{n-1} via the exponential map. Also consider the induced metric on Σ and let Z be the image of a small ball in the normal bundle to O in Σ at m by the exponential map. So Z is T' -invariant, contains m and is transversal to O . We will define an equivalence relation on a small neighbourhood V' of m in Σ by making the equivalence classes to be the orbits of U -action through points of Z . The quotient M' can be thought of a local symplectic reduction of M by the action of U . So M' is a Kahler manifold. By Theorem 1 we have a natural trivialization φ'' of the canonical bundle of M' . We have a structure-preserving T' -action on M' . Let μ^* be the restriction of μ to the dual Lie Algebra of T' . Then μ^* is a moment map for T' -action on M' . Also ξ descends to M' and the level sets of (μ^*, ξ) are SLag submanifolds of M' , which lift to level sets of (μ, ξ) on M .

We will investigate the level sets of (μ^*, ξ) on M' . Let $\tau : V' \rightarrow M'$ be the quotient map. Then $\tau : Z \rightarrow M'$ is a diffeomorphism. Let L'_m be a level set of (μ^*, ξ) through $\tau(m)$. We will prove that L'_m is diffeomorphic to an $(l+1)$ -dimensional cone and moreover all points on $L'_m - \tau(m)$ are regular points for the T' -action. We claim that **i) and iv)** follow from this. Indeed L_m is an orbit of the U -action on $\tau^{-1}(L'_m) \cap Z$. Thus L_m will be locally a product of a cone with a Euclidean ball. Also its singular set is of codimension $l+1 \geq 2$. Moreover L_m will contain regular points for the T^{n-1} -action. The differential of (μ, ξ) is surjective at those points, and thus we will deduce that the image of (μ, ξ) is open.

So we have a Kahler manifold M' with a trivialization φ'' of the canonical bundle and a structure-preserving $T' \simeq T^l$ -action, which preserves $m' = \tau(m) \in M'$ and induces an action of a maximal torus of $SU(l+1)$ on the tangent space at m' as in equation (1). We would like to understand the level set $\mu^*(m')$ of the moment map μ^* (which contains L'_m). To do this we introduce Equivariant Darboux coordinates in a following way : We identify a small neighbourhood of M' with a ball Y on a tangent bundle $T_{m'}M'$ via the exponential map. The action of T' will be linear on Y , and it will preserve the symplectic form ω' on Y , induced by the exponential map. We identify Y with a ball in \mathbb{C}^{l+1} . Then we will also have a standard symplectic form ω_0 on Y . Moser's proof of the Darboux theorem gives a embedding ϕ of a possibly smaller ball Y' into Y , s.t. $\phi^*(\omega_0) = \omega'$. Now both ω' and ω_0 are T' -invariant, so ϕ will be T' -equivariant.

The moment map of ω^0 is $\mu^0 = (\mu_1, \dots, \mu_l)$ with $\mu_i = |z_i|^2 - |z_{l+1}|^2$. The non-regular points of the action are, as we saw, the planes $(z_i = z_j = 0)$ and the zero set of μ^0 intersects them only at the origin. The zero set of μ^0 is a cone $|z_i| = |z_j|$ in \mathbb{C}^{l+1} . It's symplectic reduction thus will be a 2-dimensional cone with a singular point at the origin. Similarly we can take a level set P_0 of μ^* through m' to get a symplectic reduction M'' . Let $m'' \in M''$ be the image of m' under the quotient map. We take a fixed compact neighbourhood K of m'' in M'' and K will be a 2-dimensional cone with boundary.

We had a holomorphic function $\eta + i\xi$ on M as before, and this function descends to a holomorphic function on M' and on M'' . W.l.o.g. we assume that $\xi(m) = 0$. The zero set of ξ on M'' lifts to the fiber L'_m . The gradients of ξ and η on M are orthogonal to the orbits of T^{n-1} -action. Thus the gradient flow of, say, η on M projects to the gradient flow of η on M' and on M'' . Also the gradients of ξ and η are linearly independent over $M'' - m''$ and $\nabla\eta = J\nabla\xi$. Thus the gradient flow of η is the Hamiltonian flow of ξ , and so it preserves ξ . Since $d\xi \neq 0$ on $M'' - m''$ then near every point on the zero set of ξ in $M'' - m''$, $\xi^{-1}(0)$ is an orbit of the gradient flow of η . The orbits of this gradient flow on $\xi^{-1}(0)$ are obviously isolated. One end of such orbit might flow to the critical point m'' , but the other must flow to the boundary of K . From this we deduce that there are only finitely many of these orbits in $\xi^{-1}(0)$ in K . We will look even for a smaller $K' \subset K$ so that one end of each orbit in K' will flow to m'' . So these will be orbits d_1, \dots, d_p . We will prove that $p > 0$. We claim that from this it follows that L'_m diffeomorphic to a cone modeled on a p (l)-tori (i.e p l -tori will be the base of the cone).

Indeed let P_0 be the level set $\mu^*(m')$ of μ^* on M' as before and let $\nu : P_0 \mapsto M''$ be the quotient map. Take a point q_i on $\nu^{-1}(d_i)$ in M' . Then the gradient flow of η through q_i projects under ν to the gradient flow on M'' . Hence the gradient flow of η through q_i must terminate at m' . Let d'_i be the trajectory of this flow. Then the orbit D_i of T' -action on d'_i is diffeomorphic to a cone modeled on an l -torus. Moreover $\nu^{-1}(d_i) = D_i$. So L'_m is diffeomorphic to a cone, modeled at p l -tori.

Finally we prove that $p > 0$. We have μ^0 and on every non-regular point of T' -action some of the functions $\mu_i - \mu_j$ vanish. The set of all such points on which some of $\mu_i - \mu_j$ vanish is a union of hypersurfaces in M' . If we take a

point s outside of these hypersurfaces then the level set P_s of the moment map μ^* containing s is smooth. Take now a sequence of such points s_j converging to m' . Fix a compact neighbourhood B of m' in M' . We consider the positive direction η -gradient flow lines a_j through s_j . There are no critical points of η on P_{s_j} . So those lines a_j must intersect the boundary ∂B . We saw that the level set of P_0 is smooth outside of m' . So P_{s_j} converge to P_0 outside of m' . It is easy to show that (after passing to a subsequence) a_j converge to a flow line on P_0 terminating at m' . We project it to M'' and get a gradient flow line on M'' terminating at m'' and we are done. Q.E.D.

Remark: The proof of Theorem 2 in fact gave a construction of Special Lagrangian submanifolds as level sets

$$\mu = a, \quad \xi = c$$

Thus we effectively got an algebraic construction of Special Lagrangian submanifolds. We will utilize this construction for some examples in Section 4.

The countable union of planes in Theorem 2 stems from the fact that M is non-compact. If we make certain assumptions on the set of non-regular points of the T -action, then we can replace the countable union by a finite union:

Theorem 3 *Let $k = n - 1$ as in Theorem 2. Suppose that the set of non-regular points of the T -action on M is a finite union $S = \bigcup S_i$ of connected submanifolds s.t. each S_i has a positive-dimensional stabilizer $T_i \subset T$. Then for all points p outside a finite union $H = \bigcup H_i$ of $(n - 2)$ -planes in \mathbb{R}^n , the fiber $(\mu, \xi)^{-1}(p)$ is a smooth SLag submanifold of M .*

Proof: We have $d\xi = \text{Im}\varphi'$. On each S_i the action is non-regular, thus $\varphi' = 0$ on S_i . In particular $d\xi|_{S_i} = 0$, i.e ξ is a constant ξ_i on S_i .

Let $0 \neq e_i$ be an element in the Lie algebra of T_i . Then the flow v.field X_i of e_i vanishes along S_i . Thus $d\mu(e_i) = i_{X_i}\omega = 0$ along S_i . So in particular $\mu(e_i) = \mu_i = \text{const.}$ So the image of μ on S_i lives on a hyperplane in the dual Lie algebra \mathcal{G}^* of T .

From all this we deduce that the image of (μ, ξ) on S lives on a finite union H of $(n - 2)$ -planes in \mathbb{R}^n . Q.E.D.

3 Group actions and coassociative submanifolds

Let M^7 be a 7-manifold with a G_2 -form φ . This means that for each point $m \in M$ there is an isomorphism σ between the tangent space $T_m M$ and \mathbb{R}^7 s.t. $\sigma^*(\varphi_0) = \varphi$, there φ_0 is the standard G_2 form on \mathbb{R}^7 (see [11]). Since the group G_2 preserves the Cayley product on $\mathbb{R}^8 = \mathbb{R}^7 \oplus \mathbb{R}$, then the bundle $TM \oplus \mathbb{R}$ over M acquires a structure of an algebra, isomorphic to Cayley numbers (see [11]).

We will assume that φ is closed and co-closed, hence parallel and the Holonomy of M is contained in the group G_2 . The 4-form $*\varphi$ is a calibration and a calibrated submanifold L is called a coassociative submanifold. This is equivalent to $\varphi|_L = 0$. We will use group actions to construct coassociative submanifolds on M . We assume that $b_1(M) = 0$. We will treat 3 cases :

1) A 3-torus action. Let v_i be a basis of the Lie Algebra of T^3 and X_i are the corresponding (commuting) flow vector fields on M . Then we define the 1-forms $\sigma_1 = i_{X_2}i_{X_3}\varphi$, $\sigma_2 = i_{X_3}i_{X_1}\varphi$, $\sigma_3 = i_{X_1}i_{X_2}\varphi$. As in section 2, one can easily show that σ_i are closed, T^3 -invariant 1-forms. Hence $\sigma_i = df_i$ for some T^3 -invariant functions f_i . Since f_1 is T^3 -invariant we have

$$0 = df_1(X_1) = \sigma_1(X_1) = \varphi(X_1, X_2, X_3)$$

Consider now a level set $L = (f_i = \text{const})$. Suppose some point $m \in L$ is a regular point of T^3 -action. Since $\varphi(X_1, X_2, X_3) = 0$ one easily sees that σ_i are linearly independent at m and hence m is a smooth point of L . Also L is T^3 -invariant, hence $X_i(m)$ are in the tangent bundle of L at m . Hence one easily deduces that $\varphi|_L = 0$, i.e. L is coassociative.

2) A 2-torus action. We have vector fields X_1 and X_2 and a 1-form $\sigma = i_{X_1}i_{X_2}\varphi$. Once again $\sigma = df$ for a (T^2 -invariant) f . Consider a level set $N = (f = \text{const})$. Suppose T^2 acts freely on N and consider the quotient $M_{red} = N/T^2$, which we call a G_2 -reduction. We have a projection $\pi : N \mapsto M_{red}$. Consider the (closed) 2-forms $\eta_i = i_{X_i}\varphi$ and $\omega = i_{X_1}i_{X_2}(*\varphi)$. One can easily show that there are unique, closed 2-forms η'_i and ω' on M_{red} s.t. $\pi^*(\eta'_i) = \eta_i$ and $\pi^*(\omega') = \omega$. Let L' be a 2-submanifold of M_{red} and $L = \pi^{-1}(L')$. Then obviously L is coassociative iff $\eta'_i|_{L'} = 0$. We will reformulate this condition as a pseudoholomorphic condition on L' .

Consider a bundle $V = \text{span}(X_1, X_2)$ over N . Pick $n \in N$ and let e_1, e_2 be an orthonormal basis of V at n , compatible with the orientation, given by X_1 and X_2 . Let $e_3 = e_1 \times e_2$ (here \times is the Cayley product). e_3 doesn't depend on the choice of e_1 and e_2 and thus gives rise to a section of TM over N . Consider the bundle W over N , which is the orthogonal complement of $V \oplus (e_3)$ in TM . Then one easily verifies that the tangent bundle of N is $W \oplus V$ and the quotient of W by T -action can be viewed as a tangent bundle to M_{red} . Let again $n \in N$ and let J_i be a right Cayley multiplication by e_i . Then J_i preserve the fiber W_n of W at n and they give complex structures on W_n , which form a HyperKahler package. Also J_3 gives rise an almost complex structure on M_{red} . Let ω_i be the corresponding symplectic forms on W_n . Then one easily verifies that ω is proportional to ω_3 on W_n and $\text{span}(\eta_1|_{W_n}, \eta_2|_{W_n}) = \text{span}(\omega_1, \omega_2)$ in the space of 2-forms on W_n . From all this linear algebra we get that ω' is a symplectic form on M_{red} and J_3 is a compatible almost complex structure. Also for a 2-submanifold L' of M_{red} the conditions $\eta'_i|_{L'} = 0$ are equivalent to L' being J_3 -holomorphic.

3) An $SO(3)$ action. We will also assume that the action is not regular in at least one point. Let e_1, e_2, e_3 be the o.n. basis of the Lie Algebra of $SO(3)$. We have the following relations :

$$[e_1, e_2] = e_3, [e_2, e_3] = e_1, [e_3, e_1] = e_2$$

Let X_i be the corresponding vector fields on M . Let $\sigma = i_{X_1}i_{X_2}\varphi$. Then

$$d\sigma = \mathcal{L}_{X_1}(i_{X_2}\varphi) = i_{[X_1, X_2]}\varphi = i_{X_3}\varphi$$

Let $f = \varphi(X_1, X_2, X_3)$. Then

$$df = \mathcal{L}_{X_3}\sigma - i_{X_3}d\sigma$$

We easily deduce that both terms are 0, hence $df = 0$. Also $f = 0$ in at least 1 point. Hence $f = 0$. So we might hope to find coassociative submanifolds, invariant under $SO(3)$ -action.

Let $\alpha = i_{X_1}i_{X_2}i_{X_3}(*\varphi)$. Then using arguments as before we can show that α is a closed, $SO(3)$ -invariant 1-form. So $\alpha = dg$ for an $SO(3)$ -invariant function g . Let $v = \nabla g$. Let m be a regular point of the action. Then $v \neq 0$ at m . Also the scalar product of v and X_i is 0, so v and X_i are linearly independent and span a 4-dimensional space W . Using some Cayley algebra one can easily show that W is a coassociative subspace of TM .

We assume that $SO(3)$ acts freely on the space M' of regular points of the action. Also the complement $M - M'$ corresponds precisely to the critical points of g . Let l be a non-constant trajectory of the gradient flow of g . Then l is contained in M' . Let $L = SO(3) \times l$. Then L is coassociative. Also trajectories of the gradient flow are embedded 1-submanifolds, g is $SO(3)$ -invariant and increases on the trajectories. From all this we deduce that L is an embedded submanifold. Thus M' is covered by a family of non-intersecting coassociative submanifolds, diffeomorphic to $SO(3) \times \mathbb{R}$.

We can't in general say anything about the set of non-regular points. We will do this in one example in section 4.

4 Examples

In this section we will give a number of examples, there results of the two previous sections are applicable.

4.1 \mathbb{C}^n

There is a T^{n-1} action on \mathbb{C}^n given by

$$(e^{i\theta_1}, \dots, e^{i\theta_{n-1}}) \cdot (z_1, \dots, z_n) = (e^{i\theta_1}z_1, \dots, e^{i\theta_{n-1}}z_{n-1}, e^{-i(\theta_1 + \dots + \theta_{n-1})}z_n)$$

The moment map for this action is $\mu = (\mu_1, \dots, \mu_{n-1})$ with $\mu_i = |z_i|^2 - |z_n|^2$. The 1-form φ' , defined in the proof of Theorem 1, is $\varphi' = i^{n-1}\Sigma dz_i(z_1 \cdots \hat{z}_i \cdots z_n) = d(i^{n-1}z_1 \cdots z_n)$. So the fibration is given by

$$|z_i|^2 - |z_n|^2 = c_i, \quad \text{Im}(i^{n-1}z_1 \cdots z_n) = c_n$$

This is a classical example of Harvey and Lawson (see [7]).

4.2 $K(N)$ and the Calabi construction

Consider \mathbb{C}^n and a \mathbb{Z}_n -action on it with $k \in \mathbb{Z}_n$ acts by multiplication by $e^{2\pi ki/n}$. Then the quotient has a resolution of singularities, which is a total space of

$\gamma^{\otimes n}$, there γ is the universal line bundle over $\mathbb{C}P^{n-1}$, i.e. the resolution of singularities is the total space of the canonical bundle over $\mathbb{C}P^{n-1}$.

Let $K(N)$ be a total space of a canonical bundle of a complex manifold N and $\pi : K(N) \mapsto N$ be a projection. There is a canonical $(n,0)$ -form ρ on $K(N)$ defined by $\rho(a)(v_1, \dots, v_n) = a(\pi_*(v_1), \dots, \pi_*(v_n))$, $a \in K(N)$. The form $\varphi = d\rho$ is a holomorphic volume form on $K(N)$. If z_1, \dots, z_n are local coordinates on N then $(z_1, \dots, z_n, y = dz_1 \wedge \dots \wedge dz_n)$ are coordinates on $K(N)$ and $\varphi = dz_1 \wedge \dots \wedge dz_n \wedge dy$.

$\mathbb{C}P^{n-1}$ is a Kahler-Einstein manifold with positive scalar curvature. If N is a K-E manifold with positive scalar curvature then $K(N)$ has a Ricci-flat Kahler metric on it (see [12], p.108). The metric is constructed as follows : The connection on $K(N)$ induces a horizontal distribution for the projection π , with a corresponding splitting of the tangent bundle of $K(N)$ into horizontal and vertical distributions. We can view the horizontal space at each point $m \in K(N)$ as a tangent space to N at $\pi(m)$. Let $r^2 : K(N) \mapsto \mathbb{R}_+$ be the square of the length of an element in $K(N)$ and $u : \mathbb{R}_+ \mapsto \mathbb{R}_+$ be a positive function with a positive first derivative. We define the metric ω_u on $K(N)$ as follows: We put the horizontal and the vertical distributions to be orthogonal. On the horizontal distribution we define the metric to be $u(r^2)\pi^*(\omega)$ and on the vertical distribution we define it to be $t^{-1}u'(r^2)\omega'$. Here ω is the Kahler-Einstein metric on N , t is it's scalar curvature and ω' is the induced metric on linear fibers of π . The Kahler-Einstein condition ensures that the corresponding 2-form ω_u defining this metric on $K(N)$ is closed, i.e. the metric is Kahler. If we take $u(r^2) = (tr^2 + l)^{\frac{4}{n+1}}$ for some constant l (see [12], p.109), then ω_u is Ricci-flat.

For $\mathbb{C}P^{n-1}$ the Ricci-flat metric on $K(\mathbb{C}P^n)$ has a Kahler potential f outside of the zero section and f is a function $f = h(r^2)$, there $r^2 = \sum |z_i|^2$ on $(\mathbb{C}^n - 0)/\mathbb{Z}_n$. For instance then $n = 2$ we have the Eguchi-Hanson potential $h(x) = \sqrt{x^2 + 1} + \log x - \log(\sqrt{x^2 + 1} + 1)$. Also the metric is asymptotic to the Euclidean metric on $\mathbb{C}^n/\mathbb{Z}_n$ at infinity.

We have an $(n-1)$ -torus action on \mathbb{C}^n as in the first example, and this action commutes with the \mathbb{Z}_n -action, hence it induces an action on $K(\mathbb{C}P^{n-1})$. This action preserves the Calabi-Yau structure on $K(\mathbb{C}P^{n-1})$, and hence Theorem 2 applies. To compute the moment map, we note that $\omega_u = i\bar{\partial}\partial f = d(i\partial f)$. Now the T^{n-1} -action preserves f and so it preserves ∂f . Let $v \in \mathcal{G}$ and X_v be the vector field on $K(\mathbb{C}P^{n-1})$ as before. Then

$$0 = \mathcal{L}_{X_v}(i\partial f) = i_{X_v}\omega_u + d(i\partial f(X_v))$$

Now $i\partial f(X_v) = i(df(X_v) - idf(JX_v))$. Also $df(X_v) = 0$, so $i\partial f(X_v) = df(JX_v) = h'(r^2) \cdot dr^2(JX_v)$. If $v = \partial_{\theta_i}$ then $dr^2(JX_v) = |z_n|^2 - |z_i|^2$.

So the moment map $\mu = (\mu_1, \dots, \mu_{n-1})$ satisfies $d\mu_i = i_{X_i}\omega_u = d(h'(r^2) \cdot (|z_i|^2 - |z_n|^2))$. So $\mu_i = h'(r^2)(|z_i|^2 - |z_n|^2)$. By similar reasoning the function ξ is given, as in the previous example, by $\xi = Im(i^{n-1}z_1 \cdots z_n)$. So SLag fibration on $K(\mathbb{C}P^n)$ is given by

$$h'(r^2) \cdot (|z_i|^2 - |z_n|^2) = c_i, \quad Im(i^{n-1}z_1 \cdots z_n) = c_n \quad (2)$$

Also $h'(r^2)$ converges to 1 at infinity, so this fibration is asymptotic at infinity to the fibration on \mathbb{C}^n in the previous example.

We will now make the following general observation : Let L be an oriented Lagrangian submanifold of a K-E manifold N . We endow $K(N)$ with a Kahler metric ω_u as above for any choice of a function u . For any point $l \in L$ there is a unique element κ_l in the fiber of $K(N)$ over l which restricts to the volume form on L . Various κ_l give rise to a section κ of $K(N)$ over L . For any $\lambda \in \mathbb{R}$ we define a submanifold $L_\lambda \subset K(N)$ by

$$L_\lambda = \{(l, \mu) | l \in L, \mu = (a + i\lambda)\kappa_l, a \in \mathbb{R}\}$$

We have the following:

Lemma 1 *L is a minimal Lagrangian submanifold of N iff any of L_λ is a Special Lagrangian submanifold of $N(K)$*

Proof : First we note that L_λ are Special, i.e. $Im\varphi|_{L_\lambda} = 0$. Indeed one easily verifies that $Im\rho|_{L_\lambda} = \lambda\pi^*(\kappa)$ and hence $Im\varphi|_{L_\lambda} = \lambda\pi^*(d\kappa) = 0$.

We now prove that L_λ is Lagrangian iff L is minimal. Let m be a point on L_λ , $l = \pi(m)$ and $m = (a + i\lambda)\kappa_l$. The tangent space of L_λ at m is spanned by κ_l (viewed as a vertical vector in $T_m K(N)$) and vectors $(e + (a + i\lambda)\nabla_e \kappa)$. Here e is any tangent vector to L at l (viewed as an element of the horizontal distribution of $T_m K(N)$) and $(a + i\lambda)\nabla_e \kappa$ lives in the vertical distribution of $T_m K(N)$. To compute $\nabla_e \kappa$ take an orthonormal frame (v_j) of $T_l L$ and extend it to an orthonormal frame of L in a neighbourhood U of l in L s.t. $\nabla^L v_i = 0$ at l (here ∇^L is the Levi-Civita connection of L). We get that

$$\nabla_e \kappa = \kappa \cdot \nabla_e \kappa(v_1, \dots, v_n) = \kappa(e(\kappa(v_1, \dots, v_n)) - \sum \kappa(v_1, \dots, \nabla_e v_j, \dots, v_n))$$

Now $e(\kappa(v_1, \dots, v_n)) = 0$. Also clearly

$$\kappa(v_1, \dots, \nabla_e v_j, \dots, v_n) = i \langle \nabla_e v_j, Jv_j \rangle = i \langle \nabla_{v_j} e, Jv_j \rangle = -i \langle e, J(\nabla_{v_j} v_j) \rangle$$

Here J is the complex structure on N . Thus we get that

$$(a + i\lambda)\nabla_e \kappa = i(a + i\lambda)(Jh \cdot e)\kappa_l$$

Here $h = \sum \nabla_{v_j} v_j$ is the trace of the second fundamental form of L . From this one easily deduces that L_λ is Lagrangian iff $h = 0$, i.e. L is minimal. Q.E.D.

We will now investigate toric K-E manifolds. For recent results and examples we refer the reader to [2] and [14]. We begin with the following lemma.

Lemma 2 *Let (M^{2n}, ω) be a compact symplectic manifold and g some Riemannian metric on M . Suppose that we have an effective Hamiltonian n -torus action on M , which preserves g . Then there is a regular orbit of the action, which is a minimal submanifold with respect to g .*

Here by regular orbits we mean orbits with a finite stabilizer.

Proof: We have a moment map μ and smooth orbits are levels set of the moment map. For an orbit L to be a minimal submanifold, it is obviously necessary to be a critical point of the volume functional on the orbits. We note that it is also sufficient. Indeed let v be any element of the Lie Algebra \mathcal{G} of the torus T^n . Then $\mu(v)$ is T^n -invariant, and so is the gradient $\nabla\mu(v)$. Also this gradient is orthogonal to the orbits. Consider now this gradient flow. It commutes with T^n -action, hence it sends orbits to orbits. Since L is critical for the volume functional, we get from the first variation formula $\int_L h \cdot \nabla\mu(v) = 0$. Here h is a trace of the second fundamental form of L . But both h and $\nabla\mu(v)$ are T^n -invariant, hence we are integrating a constant. So $h \cdot \nabla\mu(v) = 0$ pointwise. Now v was arbitrary, hence $h = 0$.

Finally we note that at least one orbit is critical for the volume functional on the orbits. We use the following easy

Lemma 3 *Let L be a orbit with a positive dimensional stabilizer $T' \subset T$ and $x \in L$. Then for any $\epsilon > 0$ there is a neighbourhood U of x s.t. any orbit passing through U has volume $< \epsilon$.*

Indeed we can take a (unit) vector e_1 in the Lie Algebra of T' . Then the corresponding flow vector field X_1 vanishes along L . Extend e_1 to an o.n. basis e_2, \dots, e_n of the Lie algebra of T . The vector fields X_i will have uniformly bounded lengths. We choose a neighbourhood U of x in which X_1 has sufficiently small length and it is clear that volumes of orbits through U will be sufficiently small.

So now we try maximize volume among regular orbits. Let L_i be a sequence of orbits, whose volume goes to supremum s of volumes of regular orbits. Then by the previous lemma it is clear that a limiting orbit (of some subsequence) L is regular. Now the differential of the moment map on M is surjective along L and L_i are level sets of μ . It is clear that $L_i \mapsto L$ as manifolds and hence the volume of L is s and we are done. Q.E.D.

On $\mathbb{C}P^{n-1}$ we have the following T^{n-1} -invariant minimal Lagrangian torus

$$L = ((z_1, \dots, z_n) | |z_i| = |z_j|)$$

and corresponding SLag submanifolds L_λ in $K(\mathbb{C}P^{n-1})$. Those submanifolds are invariant under our T^{n-1} -action and they are in the moduli-space (2) we constructed (in fact L_0 is a submanifold, which corresponds to $c_i = 0$ in equation (2)). Moreover any other element in our moduli-space is asymptotic at infinity to L_0 . By that we mean the following : Let B be the unit ball of $K(\mathbb{C}P^{n-1})$ with respect to r^2 . L_0 is invariant under scaling of $K(\mathbb{C}P^{n-1})$ by a real number. If L is another element in our moduli-space then scaling of L by $k \in \mathbb{R}$ as k goes to infinity converges in C^∞ to L_0 on compact subsets of $B - \mathbb{C}P^{n-1}$. It turns out that analogous situation holds for any toric K-E manifold N with positive scalar curvature.

Suppose we have an effective, structure-preserving T^n -action on N . We make the following definition: We have a T^n action on N and this action induces a T^n -action on $K(N)$. Let \mathcal{G} be the Lie algebra of T , $v \in \mathcal{G}$, X_v be the flow vector

field on N and X'_v be the flow vector field on $K(N)$. So $\pi_*(X'_v) = X_v$. Let $l \in N$ and $m \in K_l = \pi^{-1}(l)$. Let $R(m)$ be the vertical part of X'_v at m . Since $R(m)$ is vertical, it can be viewed as an element of K_l . The correspondence $m \mapsto R(m)$ is a linear correspondence on K_l . Hence there is a complex number $\sigma_l(v)$ s.t. $R(m) = \sigma_l(v)m$. At a regular point l of the T -action $\sigma_l(v)$ can also be found in a following way : Take any unit length element $\xi \in K_l$. Extend ξ along the orbit of X_v to be invariant under the flow of X_v . Then one easily computes that $\sigma_l(v) = \nabla_{X_v} \xi \cdot \xi$. Since the flow of X_v is given by holomorphic isometries, ξ has unit length. Hence $\sigma_l(v)$ is purely imaginary. Also $\sigma_l(v)$ is linear in v (because $R(m)$ is given by the vertical part of the differential of the T -action at m , and this differential is a linear map from \mathcal{G} to $T_m K(N)$). Hence $i\sigma$ can be viewed as a map from N to the dual Lie algebra \mathcal{G}^* . This map is T -invariant.

Lemma 4 *For a regular orbit L of the T^n -action, L is minimal iff $\sigma|_L = 0$*

Proof: The section κ of $K(N)$ over L we defined in Lemma 1 is T^n -invariant. Also we computed that $\nabla_{X_v} \kappa \cdot \kappa = i(Jh \cdot X_v)$ for any $v \in \mathcal{G}$. From this the lemma follows Q.E.D.

Let t be the scalar curvature of N .

Lemma 5 *The map $\mu = -it^{-1}\sigma$ is a moment map for the action. The zero set of μ is precisely 1 regular orbit L .*

Proof Let $v \in \mathcal{G}$. We need to show that $d(-it^{-1}\sigma(v)) = i_{X_v}\omega$. We will do it at a smooth point p of the action. Choose any unit length element ξ of $K(N)$ over p . We can extend ξ to be a local unit length section, invariant under the X_v -flow. We have a connection 1-form $\eta(u) = \nabla_u \xi \cdot \xi$. Then η is invariant under the X_v -flow and the K-E condition says that $id\eta = t\omega$. So

$$0 = \mathcal{L}_{X_v} \eta = d(i_{X_v} \eta) + i_{X_v} d\eta = d\sigma(v) - it(i_{X_v} \omega)$$

So μ is a moment map. By Lemma 2 one of the regular orbits L is minimal, hence it lies on the zero set of μ by Lemma 4. Obviously this orbit is isolated in the zero set of μ . Now by Atiyah's result (see [1]), the zero set of the moment map is connected, hence it must coincide with L and we are done. Q.E.D.

Lemma 6 *The map $\mu' = u\pi^{-1}(\mu)$ is a moment map for the T -action on $K(N)$.*

Proof: Let $v \in \mathcal{T}$. We need to prove that $d\mu'(v) = i_{X'_v}\omega_u$.

We will study ω_u in more detail (see [12]). Let $m \in N$ be a regular point for the T^n -action and ξ a unit length element of $K(N)$ over m . We can extend ξ to be a local unit length section of $K(N)$, invariant under the flow of X_v . ξ gives rise to a connection 1-form ψ for the connection on $K(N)$ and the Einstein condition tells that $id\psi = t\omega$. The section ξ defines a complex coordinate a on $K(N)$, which is invariant under the X'_v -flow. Also the form $b = da + a\pi^*\psi$ vanishes on the horizontal distribution (see [12], p. 108). We have $r^2 = a\bar{a}$ and $u = u(r^2)$. Also the Kahler form ω_u on $K(N)$ is given by

$$\omega_u = u\pi^*\omega - it^{-1}u'b \wedge \bar{b}$$

One directly verifies that $\omega_u = d\eta$ for $\eta = it^{-1}u\pi^*\psi - it^{-1}\frac{u d\bar{a}}{\bar{a}}$. By our construction the flow of X'_v leaves η invariant. So

$$0 = \mathcal{L}_{X'_v}\eta = i_{X'_v}d\eta + d(i_{X'_v}\eta) = i_{X'_v}\omega_u + d(it^{-1}u\psi(X_v)) = i_{X'_v}\omega - d(\mu'(v))$$

Here we used the fact that $d\bar{a}(X'_v) = 0$ and $\psi(X_v) = \sigma(v)$. So μ' is a moment map and we are done. Q.E.D.

The torus action on N induces an action on $K(N)$ and by Theorem 2 we have a SLAG fibration on $K(N)$. We want to investigate the asymptotic behavior of the fibers at infinity. We will assume that the function $u = u(r^2)$, used to define the metric ω_u on $K(N)$ goes to infinity as r^2 goes to infinity (this holds e.g. for u defining the Ricci-flat metric).

Theorem 4 *$L_0 \subset K(N)$ is a fiber of the fibration arising from Theorem 2. Moreover, any other fiber is asymptotic to it at infinity.*

Proof: Let e_1, \dots, e_n be a basis for \mathcal{G} . Let X_i be the flow fields on N and X'_i be the flow fields on $K(N)$. Then $\pi_*(X'_i) = X_i$. Let ρ be an $(n, 0)$ form on $K(N)$ as before and $\varphi = d\rho$. Then ρ is T^n -invariant and we can prove, as in Theorem 1, that $i_{X'_1} \dots i_{X'_n} \varphi = d(\rho(X'_1, \dots, X'_n))$. Also for $\xi \in K(N)$, $\rho(\xi)(X'_1, \dots, X'_n) = \xi(X_1(\pi(\xi)), \dots, X_n(\pi(\xi)))$. We also have a moment map $\mu' = -it^{-1}u\pi^{-1}(\sigma) = u\pi^{-1}(\mu)$. Thus our SLAG fibers will be given by equations

$$u(|\xi|^2)\mu_i(\pi(\xi)) = c_i, \quad \text{Im}\xi(X_1(\pi(\xi)), \dots, X_n(\pi(\xi))) = c_n$$

Here $\xi \in K(N)$. The fiber L_0 corresponds to $c_j = 0$. The asymptotic behavior of the fibers follows immediately from this formula. Q.E.D.

We can ask a similar question in general, thus giving a non-compact analog of the SYZ conjecture (see [13]): Let N be a K-E manifold with positive scalar curvature. When is it true that N has a minimal Lagrangian submanifold L and $K(N)$ is fibered by SLAG subvarieties with fibers asymptotic to L_0 at infinity?

4.3 Resolutions of singularities and (Quasi)ALE spaces

Suppose that a finite subgroup G of $SU(n)$ acts on \mathbb{C}^n and we have a crepant resolution of singularities M . D. Joyce has recently constructed a (Quasi)ALE Ricci-flat Kahler metric ω on M (see [8] and [9]).

Suppose that G is Abelian. Then there is an orthonormal basis of \mathbb{C}^n , in which the action of G is given by diagonal matrices. Consider now the T^{n-1} -action on \mathbb{C}^n as in the first example (4.1). This action commutes with the G -action, hence it induces an action on M . Also by the uniqueness property of Joyce's construction, this action preserves ω . Hence Theorem 2 applies, and we have a Special Lagrangian fibration on M .

4.4 Coassociative submanifolds

Robert Bryant and Simon Salamon have constructed in [3] some examples of complete G_2 metrics. Some examples are on total space of a spin bundle over a

3-dimensional space form. Others are on total space Λ_-^2 of anti-self-dual 2-forms over a self-dual Einstein 4-manifold. Those 3 and 4-manifolds admit isometric actions by 2-tori and by $SO(3)$, and those actions induce structure-preserving actions on the corresponding G_2 -manifolds. We will treat 1 example in detail—the total space of a spin bundle over S^3 .

The spin bundle is a bundle $V = TS^3 \oplus \mathbb{R}$ —the direct sum of the tangent bundle of S^3 with a trivial bundle. Now S^3 can be viewed as a unit sphere of quaternions. There is an S^3 action on itself, given by $q(p) = qpq^{-1}$. Here the multiplication is a quaternionic multiplication. Obviously this action becomes an $S^3/\pm 1 = SO(3)$ -action.

In [3] a G_2 -structure was constructed on the total space of V . We won't reproduce the details of the construction but only mention that the fibers of the projection of V over S^3 are coassociative. We will look for coassociative submanifolds, invariant under $SO(3)$ action.

The points ± 1 are fixed by $SO(3)$ action and the fibers over these points are $SO(3)$ -invariant coassociative submanifolds $L_{\pm 1}$. Take now any point $m \in (S^3 - \pm 1)$. Then the stabilizer of $SO(3)$ action on m is a circle. Let N_m be the orthogonal complement in the tangent space $T_m S^3$ to the orbit of $SO(3)$ -action. Then $W = N_m \oplus \mathbb{R}$ is a sub-bundle of V over $(S^3 - \pm 1)$. W is invariant under $SO(3)$ -action. Let A_m be the orbit of m under $SO(3)$ -action (A_m is diffeomorphic to S^2) and L_m be the total space of W over A_m . Then one can easily show that L_m is a coassociative submanifold, invariant under $SO(3)$ -action. Also the union of all L_m and of $L_{\pm 1}$ is precisely the set of non-regular points of the action. Also $SO(3)$ acts freely on the set of regular points of the action. By the results of section 3, the set of regular points is covered by a family of non-intersecting coassociative submanifolds. So the whole V is covered by non-intersecting, coassociative submanifolds. Those are a 3-dimensional family of submanifolds, diffeomorphic to $SO(3) \times \mathbb{R}$, a 1-dimensional family of submanifolds, diffeomorphic to $S^2 \times \mathbb{R}^2$ and 2 submanifolds, diffeomorphic to \mathbb{R}^4 .

References

- [1] M. Atiyah : Convexity and commuting Hamiltonians, Bull. London Math. Soc., vol 14, no. 46, 1982
- [2] V. Batyrev and E. Selivanova : Einstein-Kahler metrics on symmetric toric Fano manifolds, J. Reine Angew Math. 512 (1999), 225-236
- [3] R. Bryant and S. Salamon : On the construction of some complete metrics with exceptional holonomy, Duke Math. Journal, vol 58, no.3, 1989
- [4] E. Goldstein : Calibrated Fibrations, math.DG/9911093
- [5] M. Gross: Special Lagrangian Fibrations 2: Geometry, alg-geom 9809072
- [6] R. Harvey : Spinors and Calibrations, Perspectives in Mathematics, vol 9

- [7] R. Harvey and H.B. Lawson : Calibrated Geometries, Acta Math. 148 (1982)
- [8] D. Joyce : Asymptotically Locally Euclidean metrics with holonomy $SU(m)$, math.AG/9905041
- [9] D. Joyce : Quasi-ALE metrics with holonomy $SU(m)$ and $Sp(m)$, math.AG/9905043
- [10] G. Lawlor : The angle criterion, Inven. Math. 95 (1989), no.2, 437-446
- [11] H.B. Lawson and L.M. Michelsohn : Spin Geometry, Princeton University Press
- [12] S. Salamon : Riemannian Geometry and Holonomy Groups, Pitman Press
- [13] A. Strominger, S.T. Yau and E. Zaslow: Mirror symmetry is T-Duality, Nucl. Phys. B476 (1996)
- [14] G. Tian : Kahler-Einstein metrics with positive scalar curvature, Inven. Math. vol 137, 1997
- [15] S.T. Yau : On the Ricci curvature of compact Kahler manifold and the complex Monge-Ampere equation 1, Comm. Pure Appl. Math 31 (1978).

Massachusetts Institute of Technology
 E-Mail : egold@math.mit.edu